

# BULLETIN OF PURE & APPLIED SCIENCES

Vol. 37 C  
(Chemistry)

**SPECIAL ISSUE**

**National Seminar on  
INSIGHTS INTO THE  
INTERDISCIPLINARY PERSPECTIVES  
OF CHEMICAL & BIOSCIENCES**



**ISSUE NO. 1**  
January - June

**2018**

**BPAS PUBLICATIONS**  
[www.bpaspublications.com](http://www.bpaspublications.com)

Print Version ISSN 0970 4620  
Online Version ISSN 2320 320X

# Bulletin of Pure and Applied Sciences

## Section C - Chemistry

### EDITORIAL BOARD

**Convenor (Hon.):**

**Prof. Irfan Zaheer Khan**, Dept. of Chemistry, University of Maiduguri, Nigeria

**Co-convenor (Hon):**

**Dr. A. Heidari**, South University (CSU), Irvine, California, USA

### National Editorial Advisory Board

**Dr. A. Gokulakrishnan**, Dept. of Biochemistry, Islamiah College, Vaniyambadi, Vellore - 635752, India.

**Dr. Arun Kumar Sharma**, Dept. of Chemistry, M. D. S. University Ajmer - 305001, India.

**Dr. M. Mubarak Ali**, Dept. of Chemistry, Chikkaiah Naicker College, Erode - 638004, India

**Dr. Shraddha Upadhyay**, Dept of Chemistry, SV Subharti University, Meerut - 250005, India

**Dr. Umesh Kumar Singh**, Dean, Faculty of Pharmacy, SV Subharti University, Meerut - 250005, India

**Prof. (Dr.) K. R. Desai**, Director, Dept. of Chemistry, Uka Tarsadia University, Surat- 394350, India.

**Prof. B.G. Unni**, Bio-Chem Div., Regional Research Lab., Jorhat - 785006, India.

**Prof. Ch. Sarala Devi**, Dept. of Chemistry, University College for Women, Koti - 500195, India.

**Prof. Dharam Prakash**, Dept. of Chemistry, Patna University, Patna - 800005

**Prof. G. Roychoudhary**, Regional Research Lab., Bhubneshwar - 751013, India.

**Prof. J.D. Joshi**, Dept. of Chemistry, S.P. University, Vallabh Vidyanagar - 388120, India.

**Prof. K. Chowddoji Rao**, Dept. of Polymer Sci. & Tech., S.K. University, Anantapur - 515003, India.

**Prof. K.R. Desai**, Dept. of Chemistry, South Gujarat University, Surat - 396445, India

**Prof. M.N. Roy**, Dept. of Chemistry, Nagpur University, Nagpur - 440010

**Prof. R. K. Dubey**, Department of Chemistry, University of Allahabad, Allahabad - 211002, India

**Prof. S. Sadasivam**, Dept. of Bio-Chemistry, T.N. Agri. University, Coimbatore - 641003, India.

**Prof. S.K. Kumbat**, Dept. of Chemistry, J.N.V. University, Jodhpur - 342011, India.

**Prof. Saroj Aggarwal**, School of Basic & Applied Sci., GGS Indraprastha Univ., Delhi - 110006, India.

### International Editorial Advisory Board

**Dr. Jehad M Yousef**, Dept. of Biochemistry, King Abdul Aziz University, Saudi Arabia.

**Prof. Adem ONAL**, Dept. of Chemistry, Gaziosmanpasa University, 60110-Tokat, Turkey.

**Prof. Ahmet SENER**, Dept. of Chemistry, Arts & Sc. Faculty, 100 Yil University, 65080, Van-Turkey.

**Prof. F.E. Okiemen**, Dept. of Chemistry, P.M.B. 1154, University of Benin, Benincity, Nigeria

**Prof. M. Aqil**, Dept. of Chemistry, University of Maiduguri, PMB - 1069, Nigeria.

**Prof. Md. A. Salam**, Dept. of Chemistry, University of Chittagong, Chittagong, Bangladesh.

**Managing Editor:** Prof. Dr. AK Sharma

**Publication Editor:** Manoj Kumar Singh

**(A Peer Reviewed and Refereed Journal)**

(This publication is included in Abstracted/Indexed coverage of selected services i.e. ProQuest, Indian Science Abstract, Indian Citation Index (ICI), Cosmo Foundation, Germany etc.)

**(UGC Approved Journal)**

---

**Bulletin of Pure & Applied Sciences** is devoted to the international advancement of organized knowledge on all aspects of Science. It provides a forum for comments by publishing original research contributions, scientific survey, case studies, book review and letter to the Editor-in-Chief. **Bulletin of Pure & Applied Sciences** is issued six monthly, i.e. June and December of every year. The Editor-in-Chief/Managing Editor/Editorial Board etc. assumes no responsibility for the statements and opinions advanced by the contributors. The editorial staff in its work of examining papers, received for possible publication is assisted, in an honorary capacity, by a large number of distinguished scientists working in various parts of the world. **Bulletin of Pure & Applied Science** is copy-righted. Manuscripts published in the Bulletin should not be reproduced without the written permission of the Editor-in-Chief.

---

### Subscription Rates:

- Institutional: INR2000/USD80

#### *Payment detail:*

Ch/Dd No.

Date of Ch/DD

Amount paid Rs./USD

1. Advance payment required by Demand Draft payable to BPAS PUBLICATIONS payable at Delhi.
2. Cancellation not allowed except for duplicate payment.
3. Agents allowed 10% discount.
4. Claim must be made within six months from issue date.

#### *Mail all orders to*

Subscription and Marketing Manager

**BPAS PUBLICATIONS**

115-RPS- DDA Flat, Mansarover Park,

Shahdara, Delhi-110032, India

Mob: +91- 9212200909

Email: editorbulletin8@gmail.com, bpaspublications@gmail.com

Website: www.bpaspublications.com

#### Annual Subscription Rates

Print version	India Rs. 2000	U.S. \$ 80 (visit; <a href="http://www.bpaspublications.com">www.bpaspublications.com</a> ) - for the year 2018
Print version	India Rs. 3000	U.S. \$ 80 (visit; <a href="http://www.bpaspublications.com">www.bpaspublications.com</a> ) - for the year 2019

For online, please visit- [www.indianjournals.com/bpaspublications](http://www.indianjournals.com/bpaspublications)

**Special Issue on:**

**National Seminar**

on

*Insights into the Interdisciplinary Perspectives of  
Chemical and Bio Sciences*

**From 26th to 28th February 2018**

**Guest Editor**

**Dr. K. Murugan**

**Principal**

**Government Arts College,**

**Thycaud, Thiruvananthapuram, Kerala 695014**



## Government Arts College

Thycaud, Thiruvananthapuram, Kerala 695014

Phone : 0471 2323040

Email: [artscollegeoffice@tvp@gmail.com](mailto:artscollegeoffice@tvp@gmail.com)

Web: [www.gaotvm.org](http://www.gaotvm.org)



Dear Sir/ Madam

Research in emerging areas and interdisciplinary/ multidisciplinary approaches are inevitable for making major breakthroughs in scientific world..I am delighted to write the foreword for the compendium of papers of the **National seminar on Insights into the Interdisciplinary Perspectives of Chemical and Bio Sciences** by **Govt. Arts College, Trivandrum, Kerala**. The papers in connection with the seminar are an enthusiastic celebration of diversity with respect to molecules, of natural origin, intricate structure and biological relevance. It is also a unique tribute to the many young researchers who are involved in chemical synthesis. On the top of that, there are innumerable data that interweave chemistry and biology in a very appealing way. Being the focus of chemical and biological research moving from structure to function, dynamic interactions between large biopolymers and small organic molecules that cause and control processes in living organisms need to be investigated with a new vision. Chemical Biology is sure to become even more important. Science is objective and it is human intelligence and endeavor that discover and create interesting new substances. Researchers and scientists have discovered or designed more than thirty million compounds/molecules which possess biological potentialities. Blending chemistry, biology and medicine through this seminar will revolutionize the concept of life. I am confident that in the future the young minds will elucidate the chemical mechanisms of cell functions, thought and even memory. It is my hope and expectation that the papers related with the theme of the National seminar will provide an experience and referenced resource for all Science professionals caring.

Thanking you

Yours truly

**Dr. K Murugan**



# Bulletin of Pure and Applied Sciences

## Section C - Chemistry

January-June 2018  
Volume 37, Number 1

### Contents

---

- Synthesis of 1,2,3-benzotriazol-1-yl-[n-(n'-arylthiocarbamoyl)] amidines** 1-5  
Akhila V.R., Sophia K. Philip, K.N. Rajasekharan
- Structural, Bonding and Spectral Analysis of an anti-malarial drug 5-(4-chloro phenyl)-6-ethyl-2,4-pyrimidinediamine** 6-11  
Y. Sheeba Sherlin, T.Vijayakumar, J.Binoy, S.D.D. Roy and V.S. Jayakumar
- Solid state synthesis of 5-anilino-3-(3,5-dimethylpyrazol-1-yl)-1,2,4-thiadiazoles** 12-19  
K.G. Krishnapriya, K.N. Rajasekharan
- Investigations of vibrational spectra and bioactivity of ethyl N-[1-(piperidin-1-ylmethyl)benzimidazol-2-yl]carbamate** 20-25  
G.P. Sheeja Mol, D. Aruldhas
- Optical and dielectric investigations of nano crystalline scheelite  $A_{0.5}B_{0.5}MoO_4$  (A=B=Ba, Sr and Ca)** 26-34  
S. Vidya, J.K. Thomas
- Photocatalytic degradation of an acidic dye and a basic dye using heterostructured photocatalyst** 35-41  
Baiju V, Dedhila Devadathan, Raveendran R
- Antibacterial studies of polyindole based metal oxide nanocomposites** 42-52  
Dedhila Devadathan, Baiju V, Raveendran R
- 2D and 3D QSAR model generation of CDK4 inhibitors** 53-63  
Divya V, Pushpa V.L., Sarithamol S, Manoj K.B
- Sensitized luminescence from novel pyrazolone based  $\beta$ -diketonate complexes of  $Eu^{3+}$**  64-69  
D.B. Ambili Raj, S.V. Manoj, B.S. Vinod
- Waste water treatment by removal of heavy metal using EDTA-functionalized chitosan-graphene oxide nanocomposites** 70-79  
Asha Radhakrishnan, Arya Sugathan, Deepa Vijayan, B. Beena
- Effect of a pyrimidine ring formation at the Linker part on the non linear optical property of a D- $\pi$ -A type chalcone; An Insilico study** 80-87  
Lakshmi CS Nair, Balachandran S, Hubert Joe I

12.	<b>Discovery and validation of potential impact of cannabinoid receptor agonists by insilico means</b> Sarithamol S, Divya V, V.L. Pushpa, K.B. Manoj	88-98
13.	<b>Synthesis, structural characterization and uv analysis of praseodymium zirconate oxides</b> S. Srijith, V. T. Kavitha, L.R. Asitha	99-106
14.	<b>Synthesis and characterization of ZnO @ zinc terephthalate metal organic framework</b> Harisankar. A, Asif. A	107-112
15.	<b>Synthesis, spectral, thermal and antibacterial studies of oxovanadium(IV) complex of an azo dye derived from 4-aminoantipyrine</b> Athira LS, Balachandran S, Harikumaran Nair ML	113-120
16.	<b>A comparative study on the antioxidant properties of curcuminoids and its rubrocurcumin analogues</b> Jeena John, Sudha Devi R and Balachandran S	121-125
17.	<b>An integrated approach on evaluation of hydrochemical parameters of riverine systems in Trivandrum urban area along with phosphate removal studies</b> Karthika V.S, Harsha Mahadevan, V. Viswadas, Vinu V. Dev, K. Anoop Krishnan	126-130
18.	<b>Synthesis and characterization of calcium ferrite nanoparticles by solution combustion method</b> Vidya VG, Revathy L	131-136
19.	<b>Synthesis, characterization and biological studies of transition metal complexes of Mn(II), Cu(II) and Zn(II) with an azo dye (E)-4-((2,6-dihydroxyphenyl) diazenyl)benzoic acid</b> S. Mini, Aravind S.S.	137-142
20.	<b>One pot synthesis of 1-(3-methyl-4H-benzo[1,4]thiazin-2-yl) ethanone and its antimicrobial properties</b> Shajeelammal, Minitha R., Reena Ravindran	143-147
21.	<b>Development of hetero metal organic frameworks of transition metal ions</b> Rani Pavithran, Lekshmi O	148-153
22.	<b>Spectral &amp; biological characteristics of a novel uranyl proton transfer salt</b> Shiji Fazil, Reena Ravindran	154-157
23.	<b>Isolation and characterization of active compounds from <i>Jasminum multiflorum</i> leaves</b> Afinisha Deepam LS, Divya S, Jisha MJ	158-162

*Address for communication:*

**Prof. Dr. A.K. Sharma,**  
**BPAS Publications**  
 115-RPS-DDA Flat, Mansarover Park, Shahdara, Delhi-110032, India  
 (Phone-011-22138726, 9212200909)  
 Email: editorbulletin8@gmail.com, bpaspublications@gmail.com  
 Website: www.bpaspublications.com; www.indianjournals.com

## Sensitized luminescence from novel pyrazolone based $\beta$ -diketonate complexes of $\text{Eu}^{3+}$

D.B. Ambili Raj<sup>1\*</sup>, S.V. Manoj<sup>1</sup>, and B.S. Vinod<sup>1</sup>

### Author Affiliations

<sup>1</sup>P.G. & Research Department of Chemistry, SreeNarayana College, Kollam 691001

### Corresponding Author

\*D.B. Ambili Raj, P.G. & Research Department of Chemistry, SreeNarayana College, Kollam 691001

E-mail: ambilrajdb@gmail.com

Received on 13<sup>th</sup> January 2018

Accepted on 26<sup>th</sup> January 2018

### Abstract

New pyrazolone based fluorinated ligand, 3-methyl 4-(2,2,3,3,3-penta fluoro-1-hydroxypropylidene)-1-phenyl-1H-pyrazol-5(4H)-one has been synthesized and well characterized. Europium-containing complexes were synthesized by adduct formation between 3-methyl 4-(2, 2, 3, 3, 3-penta fluoro-1-hydroxypropylidene)-1-phenyl-1H-pyrazol-5(4H)-one and europium (III) complex and a bidentate neutral donor 1,10-phenanthroline (phen). The emission spectra of the complexes display characteristic sharp peaks in the 575–700 nm region associated with the  $^5\text{D}_0 \rightarrow ^7\text{F}_J$  ( $J = 0-4$ ) transitions of the  $\text{Eu}^{3+}$  ion.

**Keywords:** Lanthanide complex, PL properties, pyrazolone, fluorinated ligand,  $\text{Eu}^{3+}$

## 1. INTRODUCTION

Lanthanide ions possess fascinating optical properties and are largely involved in various technological applications which are largely governed by their interaction with light. Light emitting diodes (LED's), television and computer displays, optical fibres, optical amplifiers, lasers, as well as responsive luminescent stains for biomedical analysis, medical diagnosis, and cell imaging rely heavily on lanthanide ions [1]. The two most useful lanthanides,  $\text{Eu}^{3+}$  and  $\text{Tb}^{3+}$ , have unusual spectroscopic characteristics, including longer lifetime (milliseconds), very sharp emission bands, and large Stokes shifts [2]. Unfortunately, as a consequence of the parity (Laporte) forbidden nature of the  $4f$  transitions, the direct absorption of Ln(III) cations is only very weak, and they hence have very low molar absorption coefficients (typically less than  $10 \text{ M}^{-1}\text{cm}^{-1}$ ) and hence a lower emission which limits their practical usage. In order to circumvent these lower values of extinction coefficients, the luminescent metal ion can be chelated to a chromophore-containing group which functions as an 'antenna,' by absorbing incident light and transferring this excitation to the metal ion, which can then deactivate by undergoing its typical luminescent emission. This process of 'antenna effect' will only be effective if the triplet level of the ligand is situated slightly above the accepting energy levels of the

lanthanide ion. This is because the accepting energy levels of the visible luminescent lanthanides are all situated at relatively high energy position of UV region in the spectra. Among these antenna molecules,  $\beta$ -diketone ligand is one kind of important antennae for the  $\text{Eu}^{3+}$  and  $\text{Tb}^{3+}$  ions [3].  $\beta$ -diketonates, as they are negatively charged bidentate ligands can form 3:1 complexes with metals, which are overall neutral. Neutral ligands like phenanthroline can be coordinated in addition, because the coordination number of the ions is eight. Among the widely known ligands  $\beta$ -diketone appears to meet the requirements for lanthanide luminescent purpose. The advantage of the use of  $\beta$ -diketone ligands is that they form stable complexes with  $\text{Ln}^{3+}$  cation. The  $\pi$ - $\pi^*$  transition of  $\beta$ -diketonates is intense and occurs over a significant range of wavelengths that is appropriate for sensitization of  $\text{Ln}^{3+}$  cation luminescence. The replacement of C–H bonds in a  $\beta$ -diketone by C–F bonds is significant in the design of highly luminescent lanthanide complexes concerned the efficient emission. C–F oscillators lower the vibrational energy of the ligands and decrease the energy loss caused by ligand vibration and thereby enhances the emission intensity of the lanthanide ion. Further due to the heavy-atom effect, which facilitates inter system crossing, the lanthanide-centered luminescent properties are enhanced [4].

These are of special interest in the fabrication of OLED devices because the ternary complexes with phenanthroline or bathophenanthroline have a much lower sublimation point that allows easy sublimation of these materials on OLED devices. The lowered sublimation points are caused by distortion of the symmetry in the complexes and vacuum deposition of the complexes is relatively easily achieved.

## 2. EXPERIMENTAL

### 2.1. Materials and Methods


Commercially available chemicals: Europium (III) nitrate fluoropropionate (TCL), 1,10-phenanthroline are used without further purification. Infrared spectra of all samples were performed on KBr pellets in the 4000–400  $\text{cm}^{-1}$  region with a resolution of 4  $\text{cm}^{-1}$ , by accumulating 64 scans using a Perkin–Elmer Spectrum One FT–IR Spectrometer. A Bruker 300 MHz NMR spectrometer was used to record the  $^1\text{H}$  NMR spectrum of the ligands. The excitation and emission spectra of the samples spectra were recorded on a Spex–Fluorolog FL3 22 spectrofluorimeter equipped with a double grating 0.22m Spex 1680 monochromator and a 450W Xe lamp as the excitation source operating in the front face mode.

### 2.2. Synthesis of the Ligand

#### Synthesis of 3 – methyl 4-(2,2,3,3,3-penta fluoro-1-hydroxypropylidene)-1-phenyl -1H-pyrazol-5(4H)-one (HMPP)

0.25g (1.435 mmol) 1-Phenyl-3-methyl-5-pyrazolone and 0.111g (1.498mmol) calcium hydroxide was taken in a RB flask. It is dissolved in 1,4-dioxane (10ml). Heated in a water bath at 60°C for about half an hour. It is cooled and 0.275g (1.432mmol) of ethyl pentafluoropropionate ( $\text{C}_5\text{H}_5\text{F}_5\text{O}_2$ ) was added and stirred for 6 hours. The resulting solution was quenched with water, added 2M HCl, and the solution was extracted twice with chloroform. The organic layer was dried over sodium sulphate ( $\text{Na}_2\text{SO}_4$ ), and the solvent was evaporated. The reaction mixture is then purified by chromatography on a silica gel column with mixture of chloroform (20%) and hexane (80%) as the eluent to get the yellow solid as the product. 3-methyl 4-(2,2,3,3,3-pentafluoro-1-hydroxypropylidene)-1-phenyl-1H-pyrazol-5(4H)-one (HL).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 7.26–7.86(m, 5H, phenyl); 1.73(s, 3H,  $\text{CH}_3$  of the pyrazole ring) (ppm) IR (KBr)  $\nu_{\text{max}}$ : 2929, 1628(C=O), 1597, 1503, 1358, 1213, 749  $\text{cm}^{-1}$



Scheme 1: Synthesis of the ligand


### 2.3. Synthesis of the binary complex

The  $\text{Eu(III)}$  complex with the ligand **HL** was synthesized according to the following method  
**Eu(L)<sub>3</sub>.2H<sub>2</sub>O**: An ethanolic solution of  $\text{Eu}(\text{NO}_3)_3 \cdot 6\text{H}_2\text{O}$  (0.5 mmol) was added to a solution of HL (1.5 mmol) in ethanol in presence of NaOH (1.5 mmol). Precipitation took place immediately, and the reaction mixture was stirred for 10 hrs at room temperature (Scheme 2). The product was filtered, washed with ethanol, then with water. The complex was then purified by recrystallization from acetone–water mixture, dried and stored in a desiccator.  
IR (KBr)  $\nu_{\text{max}}$ : 3391, 1612, 1577, 1484, 1400, 1232, 1154, 757  $\text{cm}^{-1}$ .

### 2.4. Synthesis of ternary complex


Synthesis route of the complex is shown in Scheme 3. The complex was prepared by stirring equimolar solutions of  $\text{Eu(L)}_3 \cdot 2\text{H}_2\text{O}$  and 1,10–phenanthroline in  $\text{CHCl}_3$  for 24 hrs at room temperature. The product was obtained after solvent evaporation and is purified by recrystallization from chloroform-hexane mixture.

**Eu(L)<sub>3</sub>phen**: IR (KBr)  $\nu_{\text{max}}$ : 3065, 1641, 1613, 1485, 1439, 1316, 1235, 1152, 1053, 757  $\text{cm}^{-1}$ .



Scheme

2. Synthesis of binary complex  $\text{Eu(L)}_3 \cdot 2\text{H}_2\text{O}$



Scheme 3: Synthesis of ternary complex  $\text{Eu(L)}_3(\text{phen})$

## 3. RESULTS AND DISCUSSIONS

### 3.1. Synthesis and characterization of the ligand

The ligand HL was prepared as described as in Scheme 1 and was characterized by  $^1\text{H}$  NMR. In the  $^1\text{H}$  NMR spectrum of the ligand, no peak corresponding to the enolic–OH has been observed. However, the absence of a peak at  $\delta$  3.4 ppm, corresponding to the methylene proton at the fourth position of the pyrazolone ring, confirms the existence of the ligands in the enolic form.

### 3.2. Structure and characterization of Europium(III) complex

The synthesis procedure for europium complexes were shown in Schemes 2 and 3. The IR spectra of the binary complex show a broad absorption in the region 3000–3500  $\text{cm}^{-1}$ , indicating the presence of solvent molecules in the complex. The existence of solvent molecules in lanthanide complexes with  $\beta$ -diketonates is well documented [5]. On the other hand, the absence of the broad band in the region 3000–3500  $\text{cm}^{-1}$  for the ternary complex, suggests that water molecule has been displaced by the bidentate ligands. The carbonyl stretching frequency of the ligands has been shifted to lower wave

numbers in all these complexes indicating the involvement of carbonyl oxygen in the complex formation with  $\text{Eu}^{3+}$  ion. Red shifts are observed in the C=N of 1,10-phenanthroline in the ternary complexes show the involvement of nitrogen in complex formation with  $\text{Eu}^{3+}$  ion in these complexes.

### 3.3. UV–Vis Spectra

UV–Vis absorption spectra of the  $\beta$ -diketone, 1,10-phenanthroline, and their corresponding Eu(III) complexes are measured in  $\text{CH}_3\text{CN}$  ( $c=2\times 10^{-5}$ ) are shown in Figure 1. The maximum absorption bands at 288 nm for the ligands **HL** is attributed to singlet–singlet  $\pi$ – $\pi^*$  enol absorption of  $\beta$ -diketonates. Furthermore, compared with the spectra of the ligands **HL**, the absorption band of the corresponding europium complexes are all red shifted by 2 nm, which is a consequence of the enlargement of the conjugate structure of ligands after coordinating to the  $\text{Eu}^{3+}$  ion. The spectral shapes of the absorption spectra of Eu(III) complexes in  $\text{CH}_3\text{CN}$  are similar to those of the corresponding ligands, indicating that the coordination of the europium ion does not significantly influence the energy of the singlet state of the  $\beta$ -diketone ligands.





Fig. 1: UV–Visible spectra of **HL** and complexes 1–2 in acetonitrile ( $c = 2 \times 10^{-5}$  M).

The molar absorption coefficients ( $\epsilon$ ) of the ligands **HL** is calculated as  $2.81 \times 10^4$  ( $288 \text{ nm}$ )  $\text{L mol}^{-1} \text{cm}^{-1}$ , revealing that the ligand has a strong ability of absorbing light. Similarly, the determined molar absorption coefficients ( $\epsilon$ ) of the complexes  $\text{Eu}(\text{L})_3(\text{H}_2\text{O})_2$  and  $\text{Eu}(\text{L})_3(\text{phen})$  were  $7.6 \times 10^4$  ( $290 \text{ nm}$ )  $\text{L mol}^{-1} \text{cm}^{-1}$ , which is about three times of those of the corresponding ligand, indicating the presence of three ligands in each complex molecule and is in good agreement.

### 3.4. PL Properties of Europium (III) Complexes 1–2

The solid state excitation and emission spectra of the europium complexes 1–2, recorded at 298 were shown in Figure 2. The excitation spectra were obtained with emission monitored at  ${}^5\text{D}_0 \rightarrow {}^7\text{F}_2$  hypersensitive transition around 612 nm. The room temperature excitation spectra of the europium complexes exhibit a broad band between 250 and 450 nm, which can be assigned to  ${}^1\pi$ – $\pi^*$  electron transition of the ligands [6–7].



**Fig. 2.** Room-temperature excitation and emission spectra of complexes 1–2 ( $\lambda_{\text{ex}} = 380 \text{ nm}$ ), emission monitored around 613 nm.

The excitation spectra also show a series of narrow bands assigned to the 4f-4f transitions from the  ${}^7\text{F}_0$  ground level to the excited states. However, these bands are less intense than those bands corresponding to the  $\text{S}_0 \rightarrow \text{S}_1$  transition of the ligands. This result gives evidence of europium luminescence sensitization through intramolecular energy transfer from the  $\beta$ -diketonates. Further, it also proves that luminescence sensitization via excitation of the ligand is much more efficient than the direct excitation of the  $\text{Eu}^{3+}$  ion absorption level [8]. The normalized photoluminescence spectra of the complexes 1–2 in solid state under the excitation wavelengths that maximize the  $\text{Eu}^{3+}$  emission intensity are shown in Figure 2. The emission spectra of the complexes display characteristic sharp peaks in the 575–700 nm region associated with the  ${}^5\text{D}_0 \rightarrow {}^7\text{F}_J$  ( $J = 0-4$ ) transitions of the  $\text{Eu}^{3+}$  ion. The five expected peaks of the  ${}^5\text{D}_0 \rightarrow {}^7\text{F}_{0,4}$  transitions are well resolved, and the hypersensitive  ${}^5\text{D}_0 \rightarrow {}^7\text{F}_2$  transition is very intense, pointing towards a highly polarizable chemical environment around the  $\text{Eu}^{3+}$  ion, and is responsible for the red emission colour of the complexes. Further, the emission spectra of complexes show only one peak for the  ${}^5\text{D}_0 \rightarrow {}^7\text{F}_0$ , indicating the presence of a single chemical environment around the  $\text{Eu}^{3+}$  ion and also show that the metal ion occupies a low-symmetry site [9]. It is also important to mention that the emission spectra of the europium complexes do not exhibit ligand centered transitions, indicating that there is an efficient intramolecular energy transfer from the ligands to the  $\text{Eu}^{3+}$  ion.

### 3.5. Energy transfer between ligands and Europium(III)

Generally, the sensitization pathway in luminescent europium complexes consist of the excitation of the ligands from the ground state to their excited singlet states, and subsequently through the intersystem crossing of the ligands to their triplet states, following the energy transfer from the triplet state of the ligand to the central ion [10]. In this process, the 4f electrons of the  $\text{Eu}^{3+}$  ion are excited to the  ${}^5\text{D}_0$  manifold ion from the ground state, finally the  $\text{Eu}^{3+}$  ion emits when the 4f electrons undergo a transition from the excited state of  ${}^5\text{D}_0$  to the ground state [11].

## 4. CONCLUSIONS

New pyrazolone based fluorinated ligand, 3-methyl 4-(2,2,3,3,3-penta fluoro-1-hydroxypropylidene)-1-phenyl-1H-pyrazol-5(4H)-one has been synthesized and well characterized. Europium-containing complexes were synthesized by adduct formation between 3-methyl 4-(2, 2, 3, 3, 3-penta fluoro-1-hydroxypropylidene)-1-phenyl-1H-pyrazol-5(4H)-one and europium (III) complex and a bidentate neutral donor 1,10-phenanthroline (phen). The emission spectra of the

complexes display characteristic sharp peaks in the 575–700 nm region associated with the  $^5D_0 \rightarrow ^7F_J$  ( $J = 0-4$ ) transitions of the  $\text{Eu}^{3+}$  ion. The sensitization mechanism for luminescent europium complexes involves a triplet pathway, in which the transfer of energy absorbed by the ligand to the  $\text{Eu}^{3+}$  ion takes place from the ligand-centered triplet excited state. The results demonstrate that the substitution of solvent molecules by bidentatephen molecule in  $\text{Eu}(\text{L})_3(\text{H}_2\text{O})_2$  greatly enhances the metal-centered luminescence.

## REFERENCES

- [1] Bünzli, J.-C. G., Chemical Society Reviews, 110, **2010**, 2729–2755.
- [2] Binnemans, K., Handbook on the physics and chemistry of rare earths, Elsevier, Amsterdam, 35, **2005**, 107.
- [3] Biju, S.; Ambili Raj, D. B.; Reddy, M. L. P.; Kariuki, B. M., Inorganic Chemistry, 45, **2006**, 10651–10660.
- [4] Carlos, L. D.; de Mello Donega, C.; Albuquerque, R. Q.; Alves Jr. S.; Menezes, J. F. S.; Malta, O. L., Molecular Physics, 101, **2003**, 1037–1045.
- [5] Eliseeva, S. V.; Kotova, O. V.; Gumy, F.; Semenov, S. N.; Kessler, V. G.; Lepnev, L. S.; Bünzli, J.-C. G.; Kuzmina, N. P., Journal of Physical Chemistry, 112, **2008**, 3614–3626.
- [6] de Sa, G. F.; Malta, O. L.; de Mello Donega, C.; Simas, A. M.; Longo, R. L.; Santa-Cruz, P. A.; da Silva Jr, E. F., Coordination Chemistry Reviews, 196, **2000**, 165–195.
- [7] Pavithran, R.; Saleesh Kumar, N. S.; Biju, S.; Reddy, M. L. P.; Alves, Jr. S.; Freire, R. O., Inorganic Chemistry, 45, **2006**, 2184–2192.
- [8] Latva, M.; Takalo, H.; Mikkala, V. M.; Matachescu, C.; Rodriguez-Ubis, J. C.; Kanakare, J., Journal of Luminescence, 75, **1997**, 149–169.
- [9] Kai, J.; Parra, F. D.; Brito, H. F., Journal of Materials Chemistry, 18, **2008**, 4549–4554.
- [10] Xu, H.; Wang, L.-H.; Zhu, X.-H.; Yin, K.; Zhong, G.-Y.; Hou X.-Y.; Huang, W., Journal of Physical Chemistry B, 110, **2006**, 3023–3029.
- [11] Huang, C., Science Press: Beijing, **1997**.